

On page 35, line 29, please replace "Example 7.3.," with --Example 7.2.,--

On page 46, line 11, please replace "7.3." with --7.2.--

**IN THE CLAIMS**

Please cancel claims 4, 5, 22, and 23; amend claims 9, 24-26, 35, 41-48, 53, 54, 55, 59, and 62; and add four new claims 63-66.

Please amend claims 9, 24-26, 35, 41-48, 53, 54, 55, 59, and 62 as follows:

A2 3. (amended) The method of claim 1, wherein [the one or more nutrient media contain] said jasmonate-related compound or an alkyl ester thereof is added to the one or more nutrient media.

A3 9. (amended) The method of claim 8, wherein the alkyl group esterified to jasmonic acid has from one to [six] four carbon atoms.

Sub E3  
A4 24. (amended) The method of claim 18, wherein the concentration of silver ions, silver complexes, and silver-containing compounds is [1uM - 200uM] 10uM - 100uM.

A4 25. (amended) The method of claim 18, wherein the concentration of silver ions, silver complexes, and silver-containing compounds is 50[u]uM.

26. (amended) The method of claim 18, wherein the concentration of silver ions, silver complexes, and silver-containing compounds is 10[u]uM.

A5 35. (amended) The method of claim 1, claim 3, or claim 30, wherein the one or more nutrient media further comprises a polyamine.

Sub. C19  
41. (amended) The method of claim 1, wherein the said one or more taxanes recovered is at least one compound selected from the group consisting of taxol, [taxol C, 7-xylosyltaxol,] 7-epitaxol, 10-deacetyl-7-epitaxol, cephalomannine, 10-deacetyltaxol, 7-xylosyl-10-deacetyltaxol, baccatin III, and 10-deacetyl baccatin III.

42. (amended) The method of claim 1, wherein the cells are cultured in [one] a first medium, then the medium composition is changed to a second medium to induce taxane production.

Sub. C19  
43. (amended) The method of claim 42, wherein the concentration of nitrate is lower in the [production] second medium than in the [growth] first medium, and the concentration of saccharide is higher in the [production] second medium than in the [growth] first medium.

44. (amended) The method of claim 43, wherein the [growth] first medium contains nitrate at a concentration which is 2 to 10 times the nitrate concentration in the [production] second medium.

Sub. E9  
45. (amended) The method of claim 42, wherein the [production] second medium contains saccharide at a concentration which is 2 to 5 times the saccharide concentration in the [growth] first medium.

Sub. C20  
~~46. (amended) The method of claim 1, wherein the cells are cultured in media containing saccharide in a concentration of 1 - 150 g/L, and [or] nitrate ion in a concentration of 0.3 - 70 mM.~~

Sub. E10  
47. (amended) The method of claim 43, wherein the [growth] first medium contains saccharide in the concentration of 1 - 30 g/L, and [or] nitrate ion in the concentration of 2.5 - 70 mM; and the [production] second medium contains saccharide in the concentration of 4 - 150 g/L, and nitrate ion in the concentration of 0.3 - 18 mM.

48. (amended) The method of claim 43, wherein the [growth] first medium contains saccharide in the concentration of 5 - 15 g/L, and[/or] nitrate ion in the concentration of 20 - 30 mM; and the [production] second medium contains saccharide in the concentration of 35 - 55 g/L, and nitrate ion in the concentration of 2 - 7 mM.

53. (amended) The method of claim 1, claim 3, or claim 30, wherein cells of said *Taxus* species are cultivated by a fed-batch process.

54. (amended) The method of claim [27] 53, wherein the culture medium is replenished during cultivation by [fed batch culture] periodic nutrient medium exchange.

55. (amended) The method of claim 1 or claim 30, further comprising the periodic removal of said at least one or more taxanes from the nutrient media .

59. (amended) The method of claim 1 or claim 3, wherein the cells are cultured in the presence of 0.3% to 8% [w/v] v/v of carbon dioxide in the gas phase in equilibrium with the culture medium.

62. (amended) The method of claim 1 or claim 3, wherein the cells are cultured in the presence of controlled oxygen concentration between 25% to 95% of air saturation.

Please add four new claims 63-66 as follows:

63. (new) The method of claim 42, wherein the second medium comprises a jasmonate-related compound or an alkyl ester thereof.

64. (new) The method of claim 1 or claim 30, wherein a jasmonate-related compound or an alkyl ester thereof is added continuously to the cell culture.



### AMENDED CLAIMS

1. A method for producing one or more taxanes in high yields in cell culture of a *Taxus* species comprising: cultivating in suspension culture, in one or more nutrient media under growth and product formation conditions, cells of a *Taxus* species derived from callus or suspension cultures, and recovering said one or more taxanes from said cells, said medium of said cell culture, or both, wherein at least one of the one or more nutrient media comprises an enhancement agent selected from the group consisting of (a) jasmonate-related compound or an alkyl ester thereof, (b) an antiethylene agent, and (c) an inhibitor of phenylpropanoid metabolism.

2. The method of claim 1, wherein the one or more nutrient media contain an antiethylene agent which is a silver-containing compound, or a silver complex, or a silver ion.

3. (amended) The method of claim 1, wherein [the one or more nutrient media contain] said jasmonate-related compound or an alkyl ester thereof is added to the one or more nutrient media.

4. (cancelled) The method of claim 3, wherein the jasmonate-related compounds are in a concentration from  $10^{-9}$  to  $10^{-3}$ M.

5. (cancelled) The method of claim 3, wherein the jasmonate-related compounds are in a concentration from  $10^{-6}$  to  $5 \times 10^{-4}$ M.

6. The method of claim 3, wherein the jasmonate-related compounds are in a concentration from  $10^{-5}$  to  $2 \times 10^{-4}$ M.

7. The method of claim 3, wherein the jasmonate-related compound is at least one compound selected from the group consisting of jasmonic acid and its alkyl esters, dihydrojasmonic acid and its alkyl esters, and related derivatives and analogs.

8. The method of claim 3, wherein the jasmonate-related compound is at least one compound selected from the group consisting of jasmonic acid and alkyl esters of jasmonic acid.

9. (amended) The method of claim 8, wherein the alkyl group esterified to jasmonic acid has from one to [six] four carbon atoms.

10. The method of claim 8, wherein the alkyl group esterified to jasmonic acid has one carbon atom.

11. The method of claim 3, wherein the cells are cultured in the presence of heavy metal ions, heavy metal complexes, or heavy metal-containing compounds.

12. The method of claim 11, wherein the heavy metal is cobalt.

13. The method of claim 3, further wherein the cells are cultured in the presence of an antiethylene agent.

14. The method of claim 13, wherein the antiethylene agent is an ethylene-biosynthesis antagonist.

15. The method of claim 14, wherein the ethylene-biosynthesis antagonist is a compound which inhibits ACC synthase, ACC oxidase, or ethylene oxidase.

16. The method of claim 14, wherein the ethylene-biosynthesis antagonist is acetylsalicylic acid or aminooxyacetic acid.

17. The method of claim 13, wherein the antiethylene agent is an ethylene-action antagonist.

18. The method of claim 17, wherein the ethylene-action antagonist is a silver-containing compound, a silver complex or silver ion.

19. The method of claim 18, wherein the silver is at least one compound selected from the group consisting of silver thiosulfate, silver chloride, and silver oxide.

20. The method of claim 18, wherein the silver is at least one compound selected from the group consisting of silver phosphate, silver benzoate, toluenesulfonic acid silver salt, silver acetate, silver nitrate, and silver sulfate.

21. The method of claim 18, wherein the silver is at least one compound selected from the group consisting of silver pentafluoropropionate, silver cyanate, lactic acid silver salt, silver hexafluorophosphate, citric acid trisilver salt, and silver nitrite.

22. (cancelled) The method of claim 18, wherein the concentration of silver ions, silver complexes, and silver-containing compounds is 10nM - 900uM.

23. (cancelled) The method of claim 18, wherein the concentration of silver ions, silver complexes, and silver-containing compounds is 100nM - 500uM.

24. (amended) The method of claim 18, wherein the concentration of silver ions, silver complexes, and silver-containing compounds is [1uM - 200uM] 10uM - 100uM.

25. (amended) The method of claim 18, wherein the concentration of silver ions, silver complexes, and silver-containing compounds is 50[u]uM.

26. (amended) The method of claim 18, wherein the concentration of silver ions, silver complexes, and silver-containing compounds is 10[u]uM.

27. The method of claim 18, wherein the molar ratio of silver to jasmonic acid in the one or more nutrient media is less than 9.5.

28. The method of claim 1, wherein the one or more nutrient media contain an inhibitor of phenylpropanoid metabolism.

29. The method of claim 28, wherein the inhibitor of phenylpropanoid metabolism is selected from the group consisting of 3,4,-methylenedioxynitrocinnamic acid, 3,4,-methylenedioxycinnamic acid, 3,4,-methylenedioxy-phenylpropionic acid, 3,4,-methylenedioxyphenylacetic acid, 3,4,-methylenedioxybenzoic acid, 3,4,-trans-dimethoxycinnamic acid, 4-hydroxycinnamic acid, phenylpropionic acid, fluorophenylalanine, 1-aminobenzotriazole, 2-hydroxy-4,6-dimethoxybenzoic acid, SKF-525A, ammonium oxalate, vinylimidazole, diethyldithiocarbamic acid, and sinapic acid.

30. The method of claim 1, wherein the one or more nutrient media contain at least one enhancement agent selected from each of at least two of the following classes of enhancement agents: (a) jasmonic acid or an alkyl ester thereof, (b) antiethylene agents, and (c) inhibitors of phenylpropanoid metabolism.

31. The method of claim 30, wherein the jasmonic acid alkyl ester is methyl jasmonate.

32. The method of claim 1 or claim 30, wherein the one or more nutrient media further comprise an auxin-related growth regulator.

33. The method of claim 30, wherein the antiethylene agent is a silver-containing compound, a silver complex or silver ion.

34. The method of claim 30, wherein the inhibitor of phenylpropanoid metabolism is selected from the group consisting of 3,4,-methylenedioxynitrocinnamic acid, 3,4,-methylenedioxycinnamic acid, 3,4,-methylenedioxy-phenylpropionic acid, 3,4,-methylenedioxyphenylacetic acid, 3,4,-methylenedioxybenzoic acid, 3,4,-trans-dimethoxycinnamic

acid, 4-hydroxycinnamic acid, phenylpropionic acid, fluorophenylalanine, 1-aminobenzotriazole, 2-hydroxy-4,6-dimethoxybenzoic acid, SKF-525A, ammonium oxalate, vinylimidazole, diethyldithiocarbamic acid, and sinapic acid.

35. (amended) The method of claim 1, claim 3, or claim 30, wherein the one or more nutrient media further comprises a polyamine.

36. The method of claim 35, wherein the polyamine is selected from the group consisting of spermine, spermidine, putrescine, cadaverine, and diaminopropane.

37. The method of claim 1 or claim 30, wherein the one or more nutrient media further comprise a taxane precursor.

38. The method of claim 32, wherein the auxin-related growth regulator is picloram, indoleacetic acid, 1-naphthaleneacetic acid, indolebutyric acid, 2,4-dichlorophenoxyacetic acid, 3,7-dichloro-8-quinolinecarboxylic acid, or 3,6-dichloro-o-anisic acid.

39. The method of claim 1, wherein the amount of said one or more taxanes recovered is at least 3-fold greater than the amount obtained from uninduced suspension culture.

40. The method of claim 1, wherein the amount of said one or more taxanes recovered is at least 5-fold greater than the amount obtained from uninduced suspension culture.

41. (amended) The method of claim 1, wherein the said one or more taxanes recovered is at least one compound selected from the group consisting of taxol, [taxol C, 7-xylosyltaxol,] 7-epitaxol, 10-deacetyl-7-epitaxol, cephalomannine, 10-deacetyltaxol, 7-xylosyl-10-deacetyltaxol, baccatin III, and 10-deacetylbaccatin III.

42. (amended) The method of claim 1, wherein the cells are cultured in [one] a first medium, then the medium composition is changed to a second medium to induce taxane production.

43. (amended) The method of claim 42, wherein the concentration of nitrate is lower in the [production] second medium than in the [growth] first medium, and the concentration of saccharide is higher in the [production] second medium than in the [growth] first medium.

44. (amended) The method of claim 43, wherein the [growth] first medium contains nitrate at a concentration which is 2 to 10 times the nitrate concentration in the [production] second medium.

45. (amended) The method of claim 42, wherein the [production] second medium contains saccharide at a concentration which is 2 to 5 times the saccharide concentration in the [growth] first medium.

46. (amended) The method of claim 1, wherein the cells are cultured in media containing saccharide in a concentration of 1 - 150 g/L, and[/or] nitrate ion in a concentration of 0.3 - 70mM.

47. (amended) The method of claim 43, wherein the [growth] first medium contains saccharide in the concentration of 1 - 30 g/L, and[/or] nitrate ion in the concentration of 2.5 - 70 mM; and the [production] second medium contains saccharide in the concentration of 4 - 150 g/L, and nitrate ion in the concentration of 0.3 - 18 mM.

48. (amended) The method of claim 43, wherein the [growth] first medium contains saccharide in the concentration of 5 - 15 g/L, and[/or] nitrate ion in the concentration of 20 - 30 mM; and the [production] second medium contains saccharide in the concentration of 35 - 55 g/L, and nitrate ion in the concentration of 2 - 7 mM.

49. The method of claim 42, further comprising exchanging nutrient medium at least once during taxane production.

50. The method of claim 1 or claim 30, wherein said step of cultivating further comprises exchanging nutrient medium at least once during the cultivation step.

51. The method of claim 1 or claim 30, wherein nutrient medium is the same for cell culture growth and for taxane production.

52. The method of claim 1 or claim 30, wherein cells of said *Taxus* species are cultivated by a continuous or semi-continuous process.

53. (amended) The method of claim 1, claim 3, or claim 30, wherein cells of said *Taxus* species are cultivated by a fed-batch process.

54. (amended) The method of claim [27] 53, wherein the culture medium is replenished during cultivation by [fed batch culture] periodic nutrient medium exchange.

55. (amended) The method of claim 1 or claim 30, further comprising the periodic removal of said at least one or more taxanes from the nutrient media.

56. The method of claim 1 or claim 30, wherein the *Taxus* species is selected from the group consisting of *T. canadensis*, *T. chinensis*, *T. cuspidata*, *T. baccata*, *T. globosa*, *T. floridana*, *T. wallichiana*, and *T. media*.

57. The method of claim 3 or claim 30, wherein the *Taxus* species is *Taxus brevifolia*.

58. The method of claim 1, wherein the cells are cultured in the presence of 0.03% to 15% v/v of carbon dioxide in the gas phase in equilibrium with the culture medium.

59. (amended) The method of claim 1 or claim 3, wherein the cells are cultured in the presence of 0.3% to 8% [w/v] v/v of carbon dioxide in the gas phase in equilibrium with the culture medium.

60. The method of claim 1, wherein the cells are cultured in the presence of controlled oxygen concentration between 1% to 200% of air saturation.

61. The method of claim 1, wherein the cells are cultured in the presence of controlled oxygen concentration between 10% to 100% of air saturation.

62. (amended) The method of claim 1 or claim 3, wherein the cells are cultured in the presence of controlled oxygen concentration between 25% to 95% of air saturation.

63. (new) The method of claim 42, wherein the second medium comprises a jasmonate-related compound or an alkyl ester thereof.

64. (new) The method of claim 1 or claim 30, wherein a jasmonate-related compound or an alkyl ester thereof is added continuously to the cell culture.

65. (new) The method of claim 1 or claim 30, wherein the one or more nutrient media contain glutamine.

66. (new) The method of claim 3, wherein the cells are cultured in media containing saccharide in a concentration of 1 - 150 g/L, and/or nitrate ion in a concentration of 0.3 - 70mM.